

The Averaging Myth

Why One size fits all, Fits No one

In Cellular Science.



By: Bhekisizwe Thomo

Averaging Problem, Single-cell Analysis,
Macrophages,

Immunology, Research, Science
Communication

Analogy

Imagine this: You are listening to your favorite song being sung by a high school choir, the overall average sound is perfect and harmonious to your ears. But what if you could be the choir's conductor, giving each singer the microphone to sing their parts solo. You might suddenly realize that half the singers are off-key, a quarter of them are singing a different song, and a few aren't singing at all. When you listen to the group as a single collective, their individual voices blend into one "average" sound. The average might be beautiful, but it completely masks what each person is actually doing.



In modern biology we face the paradoxical dilemma known as the "Averaging Problem." For decades, researchers have studied our frontline immune system's defenders known as macrophages by collecting millions of them in a petri dish, grinding them up in a biological blender, and measuring the resulting "average." We have long analyzed and scrutinized this bulk data to understand how our immune system fights off infections. But this approach has corrupted our systems biology understanding, leading us to believe that all cells act uniformly within a given system or environment.

Is the Averaging Problem Significantly Misleading?

When studying how the immune system responds to infection, we must remember that the infection model used in the lab petri dishes is rarely 100% efficient. For instance, consider an experiment using a multiplicity of infection (MOI) of 10. In theory, if you have 1 million macrophages in a petri dish, you would add 10 million pathogens (like viruses or bacteria) and hope that each single macrophage becomes infected. But that is rarely the case, as demonstrated by landmark single-cell studies, such as the highly cited 2015

paper by Roi Avraham and colleagues in the journal Cell (Avraham, et al., 2015). Which illustrated single-cell infection dynamics to be highly heterogeneous.

Out of one million macrophage cells, usually a third will become infected with the pathogen of study. The remaining majority becomes the uninfected “bystander” cells. When researchers use the traditional bulk approach to lyse the entire plate, they are blending this heterogeneous mixture together and causing the infection cell-specific panic signals to be diluted and masked by the uninfected bystander cells' calm background signals.

Methods

To answer this question, the researchers had to abandon the traditional bulk lysis and hone in on a single-cell level. In one study, researchers developed SCoPE2, a miniaturized automated sample preparation method that allows scientists to quantitatively analyze the proteins inside single, individualized macrophages.




Figure 1: Conceptual workflow of Single-Cell Proteomics by Mass Spectrometry (SCoPE2). Adapted from Specht et al. (2021), Genome Biology (CC BY 4.0).

This technique substantially increases quantitative accuracy and throughput while lowering costs, enabling researchers to measure thousands of proteins in single, individualized macrophage cells as they grew in an environment completely free of polarizing cytokines.



In a parallel study, another team utilized a specialized microfluidic droplet approach. This allowed them to isolate individual macrophages into tiny liquid droplets to observe their specific single-cell activation response to inflammatory triggers from pathogens such as lipopolysaccharides (LPS).




Figure 3: Schematic workflow demonstrating the isolation and encapsulation of primary human macrophages into single-cell microfluidic droplets for pro-inflammatory stimulation. Adapted from Tiemeijer et al. (2023), *Frontiers in Immunology* (CC BY).

Key Findings




Figure 3C: Single-cell droplet analysis reveals that while 100% of macrophages secrete pro-inflammatory TNF α , only a small fraction secrete anti-inflammatory IL-10. Adapted from Tiemeijer et al. (2023).

At a population level, it is widely understood that macrophages produce a delicate balance of inflammatory signals (like TNF α) followed by anti-inflammatory signals (like IL-10). However, the single-cell droplet analysis revealed that this “balance” is an illusion. Exposure to LPS showed that only a small fraction of the individual macrophages actually produce IL-10 in addition to TNF- α . The average population-wide response hid the fact that most individual cells were not performing both functions.




Figure 4: A continuum of cellular states. Even when grown in the exact same environment, single-cell analysis shows that individual macrophages display a wide, continuous gradient of different protein profiles. Adapted from Specht et al. (2021), Genome Biology (CC BY 4.0).

Traditional biology assumes that cells kept in the exact same environment will behave in the exact manner. However, single-cell proteomic analysis uncovered that even in a completely homogeneous environment, macrophage proteomes are highly heterogeneous. The data revealed a continuous gradient of different protein states among the macrophages. This means that no two macrophages are exactly alike, and cellular heterogeneity emerges even without specific polarizing signals telling the cells to change.

Conclusion

These studies highlight that analyzing the bulk population of cells creates an average that actively masks the true, highly localized responses happening at the microscopic level. However, this does not mean we should discard the biological blender entirely. Traditional bulk analysis remains absolutely paramount for understanding the broader tissue microenvironment and the macroscopic interactions that keep us healthy. Relying solely on bulk analysis

is like trying to understand a complex choir orchestra by only listening to a single blended chord. It gives you the overall harmony of the concert hall, but it misses the individual performers.

Conversely, relying solely on single-cell analysis introduces a massive level of data complexity that can be incredibly difficult to interpret. If we give every single cell a microphone, the sheer volume of biological noise can become overwhelming. The future of immunology lies in synergy. By incorporating high-resolution single-cell analysis alongside traditional bulk methods, we can map both the precise individual voices and the overarching harmony they create. The perfectly average macrophage simply does not exist.




Figure 5: Extracellular control and complexity of IL-10 secretion by macrophages, demonstrating the intricate single-cell signaling networks mapped by advance analysis. Adapted from Tiemeijer et al. (2023).

Key Takeways

- The average is an illusion: Bulk data assumes uniformity, but immune responses are driven by highly distinct subsets of cells reacting differently to their environment.
- Resolution matters: By utilizing single-cell analysis and physically separating distinct cellular populations, we can finally map out the precise biological noise and spatial differences that dictate our immune responses.
- A synergistic approach: Whether we are trying to treat diseases, engineer tissues, or understand immune evasion, we must stop treating cells as a monolith while respecting the broader microenvironment.

After all, a choir with only one voice is just a solo.

References

1. Avraham, R., Haseley, N., Brown, D., Penaranda, C., Jijon, H. B., Trombetta, J. J., Satija, R., Shalek, A. K., Xavier, R. J., Regev, A., & Hung, D. T. (2015). Pathogen Record of the Host-Cell Response Reveals Macrophage Heterogeneity. *Cell*, 162(6), 1309–1321. <https://doi.org/10.1016/j.cell.2015.08.017>
2. Specht, H., Emmott, E., Petelski, A. A., Huffman, R. G., Perlman, D. H., Serra, M., Kharchenko, P., Koller, A., & Slavov, N. (2021). Single-cell proteomic and transcriptomic analysis of macrophage heterogeneity using SCoPE2. *Genome Biology*, 22(1), 50. <https://doi.org/10.1186/s13059-021-02267-5>
3. Tiemeijer, B. M., Heester, S., Sturtewagen, A. Y. W., Smits, A. I. P. M., & Tel, J. (2023). Single-cell analysis reveals TLR-induced macrophage heterogeneity, and quorum sensing dictate population wide anti-inflammatory feedback in response to LPS. *Frontiers in Immunology*, 14, 1135223. <https://doi.org/10.3389/fimmu.2023.1135223>